

AndroFeme[®] 1

Cream containing 1% w/v (10mg/mL) Testosterone B.P (17- β - Hydroxyandrost-4-en-3-one)



CONSUMER MEDICINE INFORMATION

What is in this leaflet?

This leaflet answers some of the common questions about ANDROFEME[®]. It does not contain all the available information. It does not take the place of talking to your doctor or pharmacist. All medicines have risks and benefits.

Your doctor has weighed the possible risks of you using ANDROFEME[®] against the benefits it can have for you. If you have any concerns about using ANDROFEME[®] talk to your doctor or pharmacist. Keep this leaflet with the medicine. You may need to read it again.

What ANDROFEME[®] is used for?

ANDROFEME[®] contains the active ingredient testosterone. This form of testosterone is identical to the testosterone produced by the ovaries of women and the testes of men. Testosterone plays an important role in sexual arousal, sexual response, libido, maintenance of bone, energy levels and well-being in men and women. When women suffer from a deficiency of testosterone, one or more of these areas may be affected and ANDROFEME[®] provides testosterone to address this deficiency. The skin readily absorbs testosterone and ANDROFEME[®] is a simple and effective means of getting testosterone into the bloodstream.

Prior to using ANDROFEME[®]

ANDROFEME[®] is only for use in women. ANDROFEME[®] cream should not be used if you are allergic to testosterone, ANDROFEME[®] or any of the ingredients contained within ANDROFEME[®]. These are listed at the end of this leaflet. This product contains almond oil. Do not use ANDROFEME[®] if you have cancer especially breast cancer, kidney disease, are pregnant, lactating or have high calcium levels in the blood. If you have heart disease, high blood pressure or liver disease make sure your doctor is aware of these conditions before use as ANDROFEME[®] may not be appropriate for you. Check with your doctor or pharmacist if you are unsure about whether you have any of these conditions before starting ANDROFEME[®]. If you are taking other medications check with your doctor or pharmacist before starting ANDROFEME[®]. High level athletes need to be aware of the rules governing androgen use if prescribed ANDROFEME[®] cream. Do not use ANDROFEME[®] after the expiry date which is printed on the base (crimp) of the tube. If you use this medicine after the expiry date has passed, it may not work as well. Do not use ANDROFEME[®] if the foil seal at the top of the tube is damaged or broken. ANDROFEME[®] is not indicated for use in children.

How to use ANDROFEME[®]

Your doctor will explain how to apply ANDROFEME[®]. Follow all directions exactly as they are explained. If you are unclear talk to your doctor. The dose of ANDROFEME[®] will be determined for you by your doctor. This dose is specific to your condition and should not be varied unless directed to do so by your doctor. You will need to return to your doctor at regular intervals so that the dose can be checked via a blood test.


Opening the tube

To open the tube remove the cap and peel off the foil seal. Dispose of the foil seal and replace the cap firmly after using the cream.

Measuring the correct dose of ANDROFEME[®] Cream


A measuring applicator (syringe style) in a sealed sleeve is enclosed in the ANDROFEME[®] box. The dose of ANDROFEME[®] is measured in millilitres. The applicator is marked with 0.5ml graduations for dosing accuracy. Your doctor will have determined which dose is appropriate for you. To measure the correct dose of cream insert the tip of the applicator into the open nozzle of ANDROFEME[®] cream so that the nozzle and the shoulder of the applicator are in contact. GENTLY

squeeze the base of the ANDROFEME[®] tube until cream reaches the open nozzle of the tube. At the same time slowly withdraw the plunger of the applicator. The cream will flow into the barrel of the applicator. Fill to the required dose. For example: a 0.5ml dose (5mg testosterone) needs the flat part of the plunger level with the 0.5ml mark. If there are any air bubbles in the measured dose fill slightly past the required dose mark then depress the plunger so that the excess cream flows back into the tube. Stop at the required dose mark. Remove the applicator from the nozzle of the tube and replace the cap firmly on the tube. Depress the plunger of the applicator containing the dose of ANDROFEME[®] directly onto the skin (inner arm or upper thigh). Massage the cream into the area until absorbed. Rinse the applicator in warm water after use and replace in box with ANDROFEME[®] testosterone cream ready for the next day's application. Each 1ml of ANDROFEME[®] contains 10mg of testosterone.



Where to apply ANDROFEME[®] cream

Always apply ANDROFEME[®] cream to clean dry flat areas of skin. The best area to use ANDROFEME[®] is the inner aspects of the forearms or the upper thighs of the legs. No perfume, deodorant or moisturising creams or gels should be used on the area because this may interfere with ANDROFEME[®] from being absorbed. Never apply the cream to broken or damaged skin. Do not use ANDROFEME[®] on the genital area. ANDROFEME[®] cream should be applied at approximately the same time each day. It is recommended that patients do not swim or shower until at least one hour after application of ANDROFEME[®] cream. Close skin contact with the area of application within an hour of application, by a partner or children should be avoided. This may result in the partner or child absorbing some testosterone through the skin contact.



If you miss a dose

If you forget to apply your cream, you should apply it as soon as you remember provided this is within 9 hours of your usual time of application.

Otherwise do not apply the cream until the next application is due. Missing a dose will not create an undue disruption to your treatment.

If you use too much (overdose)

Because of the way ANDROFEME[®] is used, an unintentional overdose is unlikely.

If you think that you or anyone else may have used too much ANDROFEME[®], immediately telephone your doctor or the local Poisons Information Centre for advice or go to the Accident and Emergency Department at your nearest hospital. Do this even if there are no signs of discomfort or poisoning.

If ANDROFEME[®] is accidentally swallowed or eaten contact a doctor.



CONSUMER MEDICINE INFORMATION (cont'd)

During ANDROFEME® use

Do's

Inform your doctor or pharmacist that you are using ANDROFEME® before starting any other prescribed medicine. Some medicines interact with other medications. Tell all doctors, dentists and pharmacists who are treating you that you are using ANDROFEME®. It is **very important** that you have a follow-up blood test to measure your testosterone levels **within three weeks** of starting ANDROFEME®. This is to confirm that your starting dose is correct and ensure your testosterone levels are within the normal range for women.

Do Not's

Do not give ANDROFEME® to anyone else, even if they have the same symptoms or condition as you.

Observe care when driving or operating machinery at the start of treatment using ANDROFEME® until you know how ANDROFEME® affects you. There is no evidence that ANDROFEME® will affect your ability to drive or to operate machinery.

Side-effects

Tell your doctor as soon as possible if you do not feel well while you are using ANDROFEME®.

All medicines can have side effects in some people.

Sometimes they are serious, most of the time they are not.

Some side effects may need medical treatment.

Tell your doctor if you notice any of the following that are troublesome or ongoing:

- Skin irritation, for example rash or slight redness and itching of the skin where the cream has been applied. Any irritation will usually disappear within a few days. If it persists contact your doctor or pharmacist.
- Acne
- Loss of head hair (male pattern baldness)
- Headache
- Nausea, sickness
- Swelling of the ankles
- Yellowing of the skin
- Increased body hair
- Weight gain
- Deepening of the voice

ANDROFEME® may stop you ovulating and affect your periods. Rarely-reported events associated with testosterone supplementation include:

- Electrolyte imbalances in the blood
- Changes to the tissue of the breast and endometrium (uterus)

Tell your doctor if you observe any other effects not listed above.

Do not be alarmed by this list of possible side-effects.

You most likely will not experience any of them.

ANDROFEME® has not been associated with these effects however because it contains testosterone there is a potential for side effects.

After using ANDROFEME®

Storage

Keep the tube sealed until time for use.

Replace cap firmly on the tube once opened.

Do not transfer the cream out of the original tube into another container.

Keep the tube in a cool dry place where the temperature stays below 25°C.

DO NOT FREEZE.

Once opened, the contents of the tube should be used within 12 weeks.

Keep medicines out of reach of children.

Disposal

If your doctor tells you to stop using ANDROFEME® ask your pharmacist how to dispose of any cream that is left over.

Product description

ANDROFEME® cream is a white, opaque odorless cream in a sealed laminated tube. The crimp at the base of the tube has the batch number and expiry date imprinted on it.

The tube is boxed together with a consumer information sheet and a graduated measuring applicator.

Ingredients

Each 50 ml tube contains 500 milligrams (500mg) of testosterone.

Each 1ml dose delivers 10mg of testosterone. Other ingredients in the cream include cetomacrogol 1000, cetostearyl alcohol, almond oil, dl- α -tocopherol acetate (Vitamin E acetate), butylated hydroxytoluene, anhydrous citric acid, triethanolamine, Carbomer 940, B&J Phenonip® and purified water.

ANDROFEME® is available in a 50 ml sealed laminated tube.

Sponsor

HormoneSolutions.com

on behalf of Lawley Pharmaceuticals Pty Ltd

Unit 2/ 15A Harrogate Street, West Leederville, 6007

Western Australia, AUSTRALIA

Postal Address

P.O. Box 1146,

West Leederville, 6901

Western Australia, AUSTRALIA

International Telephone: +61 8 9388 0096

Fax: +61 8 9388 0098

USA & Canada Toll Free Phone: 1-800-961-7813

Toll Free Fax: 1-800-961-7650

Website: www.hormonesolutions.com

Email: info@hormonesolutions.com

AUST L 169317

Date of Preparation: May 2011.

ANDROFEME® is a registered trademark.